

Radiation Res. 50: 528-538, June 1972

AFRRI SR71-11
SEPTEMBER 1971

AFRRI
SCIENTIFIC
REPORT

DEFENSE

FOR REFERENCE

Do Not Take From This Room

**ALTERATION OF INTESTINAL VASOACTIVITY
DURING THE DEVELOPMENT OF THE
GASTROINTESTINAL RADIATION
SYNDROME**

ARMED FORCES RADIOBIOLOGY RESEARCH INSTITUTE
Defense Nuclear Agency
Bethesda, Maryland

Approved for public release; distribution unlimited

AFRRI SR71-11

All aspects of investigative programs involving the use of laboratory animals sponsored by DoD components are conducted according to the principles enunciated in the "Guide for Laboratory Animal Facilities and Care", prepared by the National Academy of Sciences - National Research Council.

ALTERATION OF INTESTINAL VASOACTIVITY
DURING THE DEVELOPMENT OF THE
GASTROINTESTINAL RADIATION SYNDROME


J. KABAL
S. J. BAUM
L. J. PARKHURST



S. J. BAUM

Chairman

Experimental Pathology Department



M. I. VARON

Captain MC USN

Director

ARMED FORCES RADIOBIOLOGY RESEARCH INSTITUTE
Defense Nuclear Agency
Bethesda, Maryland

ACKNOWLEDGMENT

The authors are grateful to M. M. Graham for discussions and assistance with the instrumentation as well as the computer processing of the data and to S. G. Levin for discussions of the statistical evaluation. The editorial assistance of C. H. Poppe is gratefully acknowledged.

TABLE OF CONTENTS

	Page
Foreword (Nontechnical summary)	iii
Abstract.	iv
I. Introduction	1
II. Materials and Methods	2
III. Results	5
IV. Discussion	11
References	15

LIST OF FIGURES

Figure 1. Experimental setup	3
Figure 2. Changes in minute intervals after injections of norepinephrine and isoproterenol	6
Figure 3. Intestinal blood flow and blood pressure changes	8
Figure 4. Intestinal blood flow and blood pressure changes during gradual blood withdrawal and reinfusion	10
Figure 5. Intestinal vascular resistance changes	10

TABLE

Table I. Percentage Vasoactive Compensation of the Intestinal Resistance Vessels after the Administration of Norepinephrine	9
--	---

FOREWORD

(Nontechnical summary)

The terminal phase of the gastrointestinal radiation syndrome ends in the collapse of the blood circulation. This clinical picture resembles in many aspects shock induced by injury, hemorrhage, dehydration, etc. The initiating and maintaining deteriorating factor due to ionizing radiation injury (at the dose level of 1500 rads whole-body gamma-neutron radiation used in this study) is unquestionably the structural breakdown of the small intestine. However, there is a lack of understanding about those factors which modify and eventually cause irreversible clinical deterioration. One of these suspected factors might be the disproportionately increased constriction of blood vessels to the small intestine with a subsequent persistent low intestinal blood flow state. Indeed, prolonged low intestinal blood flow itself can cause shock and lethality.

In this study, the intestinal vasoactivity of blood vessel function and structure during the postirradiation period was investigated. The resistance vessels of the dog small intestine were challenged by sudden and gradual changes in blood pressure. There were indications that during the development of the gastrointestinal radiation syndrome the functional capacity of the intestinal blood vessels is altered. It seems justified to say that this altered integrity of the intestinal vasomotion could be the key factor inducing cardiovascular collapse

ABSTRACT

The functional vascular integrity of the small intestine in relationship to its morphological alterations during the postirradiation period was examined. Dogs were exposed to 1500 rads (midline tissue dose) of mixed gamma-neutron radiation. Hemodynamic parameters of an in situ intestinal loop were measured in response to the systemic administration of norepinephrine, isoproterenol and gradual bleeding and reinfusion. The results indicate that 48 hours postirradiation when the capillaries are virtually intact and the systemic hemodynamic parameters are not altered yet, the functional integrity of the intestinal vasculature is already significantly deteriorated. The intestinal resistance vessels of the irradiated animals did not exert "autoregulatory escape" as the controls did. After 72 hours the vasocompensation to the sudden blood pressure changes was also markedly changed. The fluctuation of the intestinal resistance values during the bleeding and reinfusion periods demonstrated neurohumoral imbalance in the postirradiation period. On the basis of these results it is postulated that the terminal cardiovascular collapse in irradiated animals might be due to the development of intestinal ischemic shock.

I. INTRODUCTION

One of the hemodynamic features in the terminal phase of the gastrointestinal radiation syndrome is the impairment of the small intestinal blood flow.^{9,10} This alteration is due to the disproportionately increased intestinal vasoconstriction commonly observed in states of hemodynamic deterioration leading to ischemia. The significance of this observation is enhanced by the fact that development of small intestinal ischemia has been connected with the irreversibility of different types of shock.^{1,7,15} Although in the early part of the postirradiation period apparently the systemic hemodynamic homeostasis is still maintained,¹ nevertheless some factors which might participate in the intestinal blood vessels' tone are already altered.

There is a general agreement that the radiation-induced loss of the epithelial layer of the small intestine is responsible for the initiation and continuation^{8,14} of the deteriorating sequence through impaired absorption, dehydration, electrolyte imbalance, etc. However, considerable controversy exists about the contribution of the altered intestinal vasomotion which induces the cardiovascular collapse and eventually leads to the terminal phase of the radiation injury.

In order to measure the intestinal vasoactivity and the functional capacity of the small intestinal circulation during the development of the gastrointestinal syndrome, an in situ dog intestinal loop preparation was chosen as the experimental model. First, the small intestinal resistance vessels were challenged by alpha and beta adrenergic agents in an attempt to alter immediately the intestinal pressure-flow relationship. Then, the intestinal blood vessels were exposed to gradual hemodynamic alterations by applying controlled blood withdrawal and reinfusion.

II. MATERIALS AND METHODS

Eighteen male beagles, 12 - 14 months old and weighing 9.5 - 11.5 kg were used in this study. The animals were divided into three groups: irradiated, with 48- and 72-hour postirradiation periods, and nonirradiated controls. Food was withheld from the dogs overnight before irradiation. Water was available ad libitum. At approximately 1 hour before irradiation the animals were placed in Lucite restraining cages and transferred to the exposure room of the AFRRI-TRIGA reactor where they received 1500 rads midline tissue dose of pulsed mixed gamma-neutron radiation. The AFRRI-TRIGA reactor and the exposure room have been described previously.² The methods as delineated by Pitchford and Thorp¹⁶ were used for dosimetry. The irradiated animals were subjected to the experimental procedures described below at 48 and 72 hours postirradiation. All animals were free to consume food and water ad libitum before the experimental procedure.

Anesthesia was induced by intravenous administration of 30 mg/kg sodium pentobarbital (Nembutal). After insertion of a tracheal tube, the abdomen was opened at the midline. Figure 1 shows the experimental setup. A section of the jejunal part of the intestine, usually weighing about 12 -15 grams, was chosen for the experiment. The mesenteric vein draining the intestinal segment was cannulated with a heparinized polyethylene tubing (Clay-Adams PE-320 i.d. 0.106", o.d. 0.138") and the venous outflow was registered by an electronic drop counter. The venous blood was collected in a beaker and its volume was controlled by an electronic leveler which turned on a peristaltic pump (Harvard Apparatus 1215) to return the blood to the animal via the right femoral vein when the preset volume of 15 ml was reached. The beaker originally

contained 15 ml of heparinized saline solution in a total dose of 5000 units. The venous outflow pressure was set continuously at zero. The wide cannula inserted into the mesenteric vein was always introduced far enough distally so that it was well inside the sealed opening thereby avoiding collapse of the vein by undue lateral pressure. A small mesenteric arterial side branch near the intestine was catheterized by a heparinized cannula (polyethylene tubing, Clay-Adams PE-50 i.d. 0.023", o.d. 0.038") and the arterial inflow pressure to the intestine was measured by a Statham pressure transducer (Model No. P23Db). During the surgical procedures great care was taken to leave the vessels and the nerves intact.




Figure 1. Experimental setup

Drying and cooling of the exposed parts of the intestine were avoided by covering them with a wet gauze and a plastic sheet which also allowed visual observation. The temperature was kept at 38°C with the aid of an infrared lamp connected through an automatic thermoregulator to a temperature sensor. The temperature of the animal was also maintained constant at 37.5°C with a heating pad.

The experimental procedures began 1 hour after anesthesia or about 30 minutes (base-line period) after completion of the surgical procedure. The following parameters were obtained (or calculated) and continuously recorded by an electronic recording system (Brush Instruments Division, Cleveland, Ohio): (1) intestinal mean arterial pressure (calculated from the diastolic blood pressure + 1/3 pulse pressure); (2) blood flow (venous outflow of the intestinal segment calculated as milliliters per minute per 100 g of intestine after excising and weighing the intestinal loop at the termination of the experiment; the mesenteric tissue was trimmed from the intestinal segments before weighing them); (3) intestinal resistance (expressed in peripheral resistance units as the ratio of the mean arterial pressure in millimeters of mercury and the blood flow in milliliters per minute per 100 g of intestine); and (4) heart rate. Histological sections of the jejunum were stained with hematoxylin-eosin. For capillary preparations, the Microfil technique (Canton Bio-Medical Products, Swarthmore, Pennsylvania) was used.

The experimental protocol consisted of the following procedures. To cause vasoconstriction an alpha adrenergic agent, norepinephrine (5 µg/kg), was injected intravenously and its direct and indirect hemodynamic effects on the intestinal segment were registered and calculated. After 10 minutes, maximal intestinal vasodilation was

induced by intravenous administration of 5 mg/kg of isoproterenol hydrochloride as a beta adrenergic agent and the intestinal hemodynamic responses were monitored again for a 10-minute period. Both drugs were obtained from Winthrop Laboratories, New York, N. Y.

In the second part of the experiment, the animals were bled at the rate of 40 ml/min until 200 ml of total volume (about 25 percent of the total blood volume) were removed. When this was achieved, the blood was reinfused immediately at the same flow rate. The rate of bleeding and reinfusion was controlled by a Harvard peristaltic pump connected to the femoral vein. While the animals were bled or reinfused the aforementioned hemodynamic parameters were recorded.

Statistical analysis of the data obtained in this study was performed utilizing a Scientific Data Systems 920 digital computer. This computer was programmed to calculate means, standard deviations and correlation coefficients by the least squares method and to perform a paired t-test analysis.

III. RESULTS

In the first part of the experiment the hemodynamic alterations of an in situ jejunal loop after single intravenous injections of the alpha and beta adrenergic agents were recorded. In Figure 2, the postinjection time sequence in 1-minute intervals for the mesenteric arterial pressure, the intestinal blood flow (venous outflow from the jejunal loop), intestinal resistance, and the heart rate changes are plotted.

There were no significant differences in the mean arterial pressure of irradiated and nonirradiated animals either initially or postinjection (Figure 2A). The initial venous outflows of the intact and 48-hour postirradiation intestinal loops were still in

the same range, however the blood flow values at 72 hours postirradiation decreased significantly (Figure 2B). After norepinephrine injection the sudden systemic blood pressure change resulted in a transient increased intestinal blood flow in all groups. Thereafter, the alpha adrenergic effect was manifested in a significantly sharp decline in the intestinal blood flow in all groups with the largest response in the 48-hour post-irradiation group. The beta adrenergic agent (isoproterenol) caused an approximately




Figure 2. Changes in minute intervals after injections of norepinephrine (NE) and isoproterenol (IP). (A) Blood pressure changes; (B) Intestinal blood flow changes; (C) Intestinal vascular resistance changes; (D) Heart rate changes.

★ indicates significant difference ($p < 0.01$).

30 percent decline of the intestinal blood flow in all groups within the first 3 minutes postinjection time. Thereafter, it increased steadily without returning to the original preinjection values.

The initial peripheral resistance values of the intestinal loops in the irradiated animals were higher than those of the normal controls (Figure 2C). After norepinephrine injection the control peripheral resistance balanced out within 2 minutes and remained stable contrary to that of the irradiated animals where the resistance continued to rise. However, even at these increased intestinal resistance values, the response to isoproterenol to reach maximum vasodilation appeared to be the same in all groups. Furthermore, the immediate change after 1 minute postinjection also showed the same steep slope of response.

The heart rate curves (Figure 2D) showed the characteristic responses to the adrenergic agents. Compensatory vagal reflex activity slowed the heart, overcoming the direct cardiac accelerator action of norepinephrine. The direct chronotropic action of isoproterenol was also well expressed in all groups. Although the irradiated animals' initial response average values were higher than the normal controls percentage-wise (taking the initial values as 100 percent), the magnitude of the chronotropic responses in all groups was similar.

When blood pressure was plotted against intestinal blood flow it was possible to gain a better insight into the intestinal vasoactivity induced by the two adrenergic agents. In Figure 3 the percent change of the average blood pressure was plotted against flow changes, grouped at minute intervals. It is important to point out that both adrenergic agents first altered the intestinal blood flow immediately by a sudden




Figure 3. Intestinal blood flow and blood pressure changes expressed in percentage of the preinjection values (before norepinephrine and isoproterenol, respectively). Each point represents the average of minute intervals during the postinjection period.

change of systemic blood pressure and only thereafter their local effects were manifested. The intestinal vessels of the intact and 48-hour postirradiation groups showed a gradual response after the first minute (Figure 3A,B) however the slope of the increased blood pressure to blood flow ratio of the 72-hour postirradiated intestines was followed immediately by a similar slope in the opposite direction without any compensation (Figure 3C). When the so-called vasoactive compensation for the first 2 minutes norepinephrine postinjection time was calculated, the values listed in Table I were obtained by using the following formula:

$$\text{vasoactive compensation} = \frac{(A-R)}{A} \times 100$$

where A (action) and R (reaction) of the intestinal blood flow changes (in percentage of the initial blood flow) were calculated for 100 mm Hg blood pressure change.

Table I. Percentage Vasoactive Compensation of the Intestinal Resistance Vessels after the Administration of Norepinephrine

Groups	Time intervals (min)	Change in blood pressure (mm Hg)	Change in blood flow (percent)	Percent blood flow change per 100 mm Hg blood pressure change	Vasoactive compensation (percent)
Control	0 - 1	+ 74.6	+ 26.4	+ 35.39	+ 75.2
	1 - 2	- 47.8	- 4.2	- 8.78	
48 hours postirradiation	0 - 1	+ 80.2	+ 50.0	+ 62.34	+ 42.4
	1 - 2	- 60.2	- 21.6	- 35.88	
72 hours postirradiation	0 - 1	+ 83.0	+ 57.4	+ 69.16	- 0.10
	1 - 2	- 59.3	- 41.3	- 69.65	

When the bleeding part of the experiment was started the slope of the blood pressure to blood flow regression lines of the irradiated animals (Figure 4A) was shifted significantly down to the vasoconstrictor side compared with the control. At these lower levels, the induced intestinal blood flow changes by the gradual decrease of blood pressure were similar or even somewhat better (less steep slope) than in the intact, nonirradiated animals. However, it is important to note that the control intestinal vessels showed reactive hyperemia during reinfusion which was completely missing in the irradiated intestinal resistance vessels (Figure 4B). When the intestinal peripheral resistance percentage changes were plotted against the blood pressure changes (Figure 5) during the blood withdrawal (Figure 5A) and reinfusion (Figure 5B) periods, it became clear that the intestinal blood vessels of the irradiated animals

demonstrated a considerable fluctuating vasoactivity, indicating a less coordinated vascular regulatory process. The fluctuation of the intestinal peripheral resistance was more apparent in the earlier postirradiation period, at 48 hours.




Figure 4. Intestinal blood flow and blood pressure changes during gradual blood withdrawal and reinfusion. Each point represents the average changes induced by subsequent 20-ml blood withdrawal or reinfusion. Diagonal line depicts an unchanged intestinal resistance; peripheral resistance unit is 1. Points above or under the line correspond to vasodilation or vasoconstriction, respectively.




Figure 5. Intestinal vascular resistance changes in percentage of the prehemorrhagic values according to the blood pressure alterations during gradual blood withdrawal and reinfusion periods. Each point represents the average percent change of the intestinal resistance induced by subsequent 20-ml blood volume alterations.

IV. DISCUSSION

In the present study, experiments were performed on intestinal segments which were not separated, had intact vasoconstrictor fibers and vagal innervation. Furthermore, to avoid metabolic shifting, the animals were not starved. Since the venous outflow pressure was set at zero, it is justifiable to say that the intestinal resistance was caused by the resistance vessels beginning morphologically at the arterioles and extending to the precapillary sphincters.

The present study is concerned with the functional capacity of the irradiated intestinal resistance vessels to respond when they are challenged by rapid or gradual alterations of the arterial blood pressure. In the first case apparently the reflexly mediated regulatory processes are involved. Sarnoff and Yamada¹⁷ described potent baroreceptors in the mesenteric vasculature. The pressure alterations cause activation of afferent nerve impulses from these baroreceptors with a resultant negative feedback effect. Mařorova¹² reported that irradiation can produce an inhibition or total reduction of the biopotentials (nerve conductivity).

Therefore, the possibility exists that the nerve conductance disturbance was manifested at 72 hours postirradiation when the intestinal resistance vessels appeared to be passively altered by the abrupt change of the arterial pressure.

Wallentin¹⁹ demonstrated that the vasoconstrictor fiber effect on the intestinal capillary resistance vessels is strongly counteracted by a local vasodilator mechanism. This causes an "autoregulatory escape" from the constrictor fiber influence, so balanced that in the steady-state phase of constrictor fiber activity the intestinal flow resistance is only modestly increased. Shehadeh et al.¹⁸ confirmed this

phenomenon with prolonged interarterial infusion of vasoactive drugs. The present data concerning the normal, control animals substantiate these observations. However, in the irradiated animals this so-called "autoregulatory escape" was not manifested after norepinephrine injection. The intestinal resistance continuously increased during the postinjection period with a concomitantly decreasing intestinal blood flow.

During the postirradiation period the increased resistance in the intestinal vessels (down to the precapillary sphincters) due to sympathetic activity or vasoactive agents might render ineffective the autonomous microcirculatory regulation of pressure and flow patterns within the exchange vessels. This phenomenon is not a specific hemodynamic feature of the gastrointestinal radiation syndrome since it certainly can be observed in different pathological states. It is well known however that the irradiated animals' blood vessels develop extreme reactivity to constrictor stimuli.¹¹

It was reported that in the dog and cat little or no increase in mesenteric resistance is usually seen following hemorrhage.⁵ The present data confirm this observation when in the second phase of the study the intestinal vasculature was exposed to gradual hemodynamic changes. During the blood withdrawal and reinfusion periods only a fine oscillation appeared in the resistance of the normal intestinal vessels. However, the vasomotor regulatory processes of the irradiated vessels show a marked fluctuation during the hemodynamic adaptation.

Zweifach²¹ and Windmueller et al.²⁰ indicate that the intestinal blood flow depends on the well organized local metabolic and/or some of the circulating humoral factors. A general change in the content of epinephrine and norepinephrine of various tissues was recently reported in irradiated animals.¹³

Goodall and Long⁶ showed that the physiological demand for epinephrine and norepinephrine is so great after irradiation that the adrenal gland is partially or completely depleted of its catecholamine; however, the biosynthesis of these hormones does not decrease; if anything, it increases. Similar observations were made about the adrenal cortex⁴ concerning production of glucocorticoids. Furthermore, Zweifach and Kivy-Rosenberg²² indicated that after whole-body irradiation, responses of terminal arterioles and precapillaries to epinephrine and norepinephrine are exaggerated.

The present data concerning the fluctuating intestinal resistance values when gradual blood withdrawal and reinfusion were performed substantiate the existence of the neurohumoral imbalance during the postirradiation period.

During blood reinfusion in control dogs there was a proportionately higher increase in the blood flow compared with the blood pressure increase (reactive hyperemia). This phenomenon was completely missing in the irradiated intestinal vessels. Instead, their tone was steadily shifted to the vasoconstriction side. It is necessary to point out, however, that during the development of the gastrointestinal syndrome the increased peripheral resistance has two other but less significant components; namely, increased viscosity and the formation of plugs in the capillaries (cell debris), besides active vasoconstriction.

Everhard et al.³ showed recently that decreased intestinal blood flow leads to a progressive vasoconstriction with a persistent refractory intestinal ischemia. This process, the altered integration of the intestinal vasomotion, might be one of the determinant factors in the syndrome. The progressive alterations of the hemodynamic regulation lead to hypotension, intestinal ischemia and the terminal cardiovascular

collapse, regardless of whether the electrolyte and water balance is maintained.

Therefore, it seems justifiable to conclude that the terminal phase of the gastrointestinal radiation syndrome resembles the pattern of intestinal ischemic shock.

REFERENCES

1. Caster, W. O., Armstrong, W. D. and Simonson, E. Changes in the cardiovascular system following total body X-irradiation. *Am. J. Physiol.* 188:169-177, 1957.
2. Dowling, J. H. Experimental determination of dose for the monkey in a reactor pulse environment. Bethesda, Maryland, Armed Forces Radiobiology Research Institute Scientific Report SR66-3, 1966.
3. Everhard, M. E., Regan, J. A., Veith, F. J. and Boley, S. J. Mesenteric vasomotor response to reduced mesenteric blood flow. *Physiologist* 13:191 (Abstract), 1970.
4. Fleming, K., Geierhaas, B., Hemsing, W. and Mannigel, U. Strahlenwirkungen und Nebennierenrinde. IV. Untersuchungen an Ratten über den 'ersten Corticosteroidanstieg' nach Ganzkörper-Röntgenbestrahlung. *Int. J. Radiation Biol.* 14: 93-105, 1968.
5. Friedman, J. J. Mesenteric circulation in hemorrhagic shock. *Circulation Res.* 9:561-565, 1961.
6. Goodall, McC. and Long, M. Effect of whole-body X-irradiation on the adrenal medulla and the hormones adrenaline and noradrenaline. *Am. J. Physiol.* 197: 1265-1270, 1959.
7. Gurd, F. N. Metabolic and functional changes in the intestine in shock. *Am. J. Surg.* 110:333-336, 1965.
8. Hager, E. B., Ferrebee, J. W. and Thomas, E. D. Damage and repair of the gastrointestinal tract after supralethal radiation. *Radiobiol. Radiother. (Berlin)* 4:1-11, 1963.
9. Janossy, Gy. Intestinal blood flow in gastrointestinal irradiation syndrome in the rat. *Acta Med. Acad. Sci. Hung.* 26(1):13-21, 1969.
10. Janossy, Gy. Cardiac output and its distribution in the terminal stage of the gastrointestinal irradiation syndrome ("radiation shock"). *Acta Med. Acad. Sci. Hung.* 26(1):23-29, 1969.
11. Kivv-Rosenberg, E. and Zweifach, B. W. Microcirculatory effect of serotonin on hemostasis following whole-body X-irradiation. *Am. J. Physiol.* 211:730-734, 1966.

12. Maïorova, N. F. Investigation of the bioelectric activity in the mesenteric nerves of the cat in the case of whole-body and local x-irradiation. *Radiobiologiya* 9: 764-766, 1969.
13. Minaev, P. F. and Musagalieva, G. M. Change in the content of adrenalin and noradrenalin in various tissues of irradiated animals. *Radiobiologiya* 9:625-628, 1969.
14. Osborne, J. W. Prevention of intestinal radiation death by removal of the irradiated intestine. *Radiation Res.* 4:541-546, 1956.
15. Palmerio, C., Zetterstrom, B., Shammash, J., Euchbaum, E., Frank, E. and Fine, J. Denervation of the abdominal viscera for the treatment of traumatic shock. *New Engl. J. Med.* 269:709-716, 1963.
16. Pitchford, T. L. and Thorp, J. W. The acute mortality response of beagles to pulsed, mixed gamma-neutron radiations. Bethesda, Maryland, Armed Forces Radiobiology Research Institute Scientific Report SR68-15, 1968.
17. Sarnoff, S. J. and Yamada, S. I. Evidence for reflex control of arterial pressure from abdominal receptors with special reference to the pancreas. *Circulation Res.* 7:325-335, 1959.
18. Shehadeh, Z., Price, W. E. and Jacobson, E. D. Effects of vasoactive agents on intestinal blood flow and motility in the dog. *Am. J. Physiol.* 216:386-392, 1969.
19. Wallentin, I. Studies on intestinal circulation. *Acta Physiol. Scand.* 69 (Supplementum 279):1-38, 1966.
20. Windmueller, H. G., Spaeth, A. E. and Ganote, C. E. Vascular perfusion of isolated rat gut: norepinephrine and glucocorticoid requirement. *Am. J. Physiol.* 218:197-204, 1970.
21. Zweifach, B. W. Vasoactive agents in the microcirculation. *Fed. Proc.* 27:1399-1402, 1968.
22. Zweifach, B. W. and Kivy-Rosenberg, E. Microcirculatory effects of whole-body X-irradiation and radiomimetic procedures. *Am. J. Physiol.* 208:492-498, 1965.

DISTRIBUTION LIST

AIR FORCE

The Surgeon General, U. S. Department of the Air Force, Washington, D. C. 20333 (1)
Executive Officer, Director of Professional Services, Office of the Surgeon General, Hq. USAF (AFMSPA) T-8,
Washington, D. C. 20333 (1)
Headquarters, U. S. Air Force (AFMSPAB), Washington, D. C. 20333 (1)
USAFSAM (SMBR), ATTN: Chief, Radiobiology Branch, Brooks AFB, Texas 78235 (1)
Air Force Weapons Laboratory, ATTN: WLIL (1), ATTN: WLRB-2 (1), Kirtland AFB, New Mexico 87117 (2)
Chief, Nuclear Medicine Department, P. O. Box 5088, USAF Hospital, Wright-Patterson AFB, Ohio 45433 (1)
Office of the Command Surgeon (ADCSG), Hq. ADC, USAF, Ent AFB, Colorado 80912 (1)

ARMY

The Surgeon General, U. S. Department of the Army, Washington, D. C. 20315 (1)
Surgeon General, ATTN: MEDDH-N, U. S. Department of the Army, Washington, D. C. 20315 (1)
USACDC CSSG, Doctrine Division, Fort Lee, Virginia 23801 (1)
CG, USCONARC, ATTN: ATUTR-TNG (NBC), Fort Monroe, Virginia 23351 (1)
Commanding Officer, U. S. Army Medical Research Laboratory, Fort Knox, Kentucky 40121 (1)
Commanding Officer, USA Nuclear Medical Research Detachment, Europe, APO New York, N. Y. 09180 (2)
Army Research Office, ATTN: Chief, Scientific Analysis Branch, Life Sciences Division, 3045 Columbia Pike,
Arlington, Virginia 22204 (1)
Division of Nuclear Medicine, Walter Reed Army Institute of Research, Walter Reed Army Medical Center
Washington, D. C. 20012 (5)
Commanding Officer, U. S. Army Environmental Hygiene Agency, ATTN: USAEHA-RP, Edgewood Arsenal, Mary-
land 21010 (1)
Commandant, U. S. Army Medical Field Service School, ATTN: MEDEW-ZNW, Fort Sam Houston, Texas 78234 (1)

NAVY

Chief, Bureau of Medicine and Surgery, U. S. Navy Department, Washington, D. C. 20390 (1)
Chief, Bureau of Medicine and Surgery, ATTN: Code 71, U. S. Navy Department, Washington, D. C. 20390 (1)
Director, Biological Sciences Division, Office of Naval Research, Washington, D. C. 20360 (1)
Commanding Officer, Naval Aerospace Medical Institute, NAMC, ATTN: Research Director, Pensacola, Florida
32512 (3)
Head, Animal Behavioral Sciences Branch, Naval Aerospace Medical Institute, Naval Aerospace Medical Center,
ATTN: Dr. John S. Thach, Jr., Pensacola, Florida 32512 (1)
Commanding Officer, U. S. Naval Hospital, ATTN: Director, REEL, NNMIC, Bethesda, Maryland 20014 (1)
Commanding Officer, Nuclear Weapons Training Center, Atlantic, Nuclear Warfare Department, Norfolk, Virginia
23511 (1)
Commanding Officer, Naval Submarine Medical Center, Naval Submarine Base, NL, ATTN: Medical Library,
Groton, Connecticut 06340 (1)
Commanding Officer, Naval Submarine Medical Center, Naval Submarine Base, NL, ATTN: Code 53, Nuclear Medi-
cine Training Division, Groton, Connecticut 06340 (1)

D. O. D.

Director, Defense Nuclear Agency, Washington, D. C. 20305 (1)
Director, Defense Nuclear Agency, ATTN: DDST, Washington, D. C. 20305 (1)
Director, Defense Nuclear Agency, ATTN: Chief, Medical Directorate, Washington, D. C. 20305 (4)
Director, Defense Nuclear Agency, ATTN: Technical Library (APTL), Washington, D. C. 20305 (2)
Commander, Field Command, Defense Nuclear Agency, ATTN: FC Technical Library, Kirtland AFB, New Mexico
87117 (1)
Director, Armed Forces Institute of Pathology, Washington, D. C. 20305 (1)
Administrator, Defense Documentation Center, Cameron Station, Bldg. 5, Alexandria, Virginia 22314 (12)

OTHER GOVERNMENT

U. S. Atomic Energy Commission, Headquarters Library, Reports Section, Mail Station G-17, Washington, D. C.
20545 (1)
U. S. Atomic Energy Commission, Division of Biology and Medicine, Washington, D. C. 20545 (1)
U. S. Atomic Energy Commission, Bethesda Technical Library, 7920 Norfolk Avenue, Bethesda, Maryland 20014 (1)

OTHER GOVERNMENT (continued)

National Aeronautics and Space Administration, ATTN: Lt. Col. Charles M. Barnes, USAF, DB-3, MSC, Houston, Texas 77058 (1)
National Aeronautics and Space Administration, Manned Spacecraft Center, ATTN: Dr. B. D. Newsom, Mail Code DA, Houston, Texas 77058 (1)
National Bureau of Standards, ATTN: Chief, Radiation Physics Division, Washington, D. C. 20234 (1)
U. S. Public Health Service, Bureau of Radiological Health, Division of Biological Effects, 12720 Twinbrook Parkway, Rockville, Maryland 20852 (1)
U. S. Public Health Service, Bureau of Radiological Health, Library, 12720 Twinbrook Parkway, Rockville, Maryland 20852 (1)
U. S. Public Health Service, Northeastern Radiological Health Laboratory, 109 Holton Street, Winchester, Massachusetts 01890 (1)
U. S. Public Health Service, Southeastern Radiological Health Laboratory, P. O. Box 61, Montgomery, Alabama 36101 (1)
U. S. Public Health Service, Southwestern Radiological Health Laboratory, P. O. Box 15027, Las Vegas, Nevada 89114 (1)

OTHER

Argonne National Laboratory, Library Services Department, Report Section Bldg. 203, RM-CE-125, 9700 South Cass Avenue, Argonne, Illinois 60440 (1)
Dr. Donald G. Baker, Radiobiology Department, Zellerbach Saroni Tumor Institute, 1600 Divisadero Street, San Francisco, California 94115 (1)
Dr. J. T. Brennan, Radiology Department, University of Pennsylvania, 3400 Spruce Street, Philadelphia, Pennsylvania 19104 (1)
Brookhaven National Laboratory, Information Division, ATTN: Research Library, Upton, Long Island, New York 11973 (2)
Dr. J. S. Burkle, Director of Nuclear Medicine, York Hospital, York, Pennsylvania 17403 (1)
Director, Radiobiology Laboratory, University of California, Davis, California 95616 (1)
University of California, Lawrence Radiation Laboratory, Library, Bldg. 50, Room 134, Berkeley, California 94720 (1)
University of California, Lawrence Radiation Laboratory, Technical Information Division Library L-3, P. O. Box 808, Livermore, California 94551 (2)
University of California, Laboratory of Nuclear Medicine and Radiation Biology, Library, 900 Veteran Avenue, Los Angeles, California 90024 (1)
Cdr. William H. Chapman, USN (Ret.), Bio-Medical Division L-523, Lawrence Radiation Laboratory, University of California, P. O. Box 808, Livermore, California 94551 (1)
Director, Collaborative Radiological Health Laboratory, Colorado State University, Fort Collins, Colorado 80521 (1)
Dr. L. W. Davis, Radiology Department, University of Pennsylvania, 3400 Spruce Street, Philadelphia, Pennsylvania 19104 (1)
Professor Merrill Eisenbud, New York University, Tuxedo, New York 10987 (1)
Dr. T. C. Evans, Radiation Research Laboratory, College of Medicine, University of Iowa, Iowa City, Iowa 52240 (1)
Dr. Arnold Feldman, Institute of Radiology, School of Medicine, Washington University, 510 South Kingshighway, St. Louis, Missouri 63110 (1)
Mr. Orin Gelderloos, Division of Literature, University of Michigan, Dearborn Campus, Dearborn, Michigan 48124 (1)
General Dynamics/Fort Worth, ATTN: Librarian, P. O. Box 748, Fort Worth, Texas 76101 (1)
Gulf General Atomic Incorporated, ATTN: Library, P. O. Box 608, San Diego, California 92112 (1)
IIT Research Institute, ATTN: Document Library, 10 West 35th Street, Chicago, Illinois 60616 (1)
Dr. R. F. Kallman, Department of Radiology, Stanford University, Palo Alto, California 94305 (1)
Dr. L. S. Kelly, Donner Laboratory, University of California at Berkeley, Berkeley, California 94720 (1)
Los Alamos Scientific Laboratory, ATTN: Report Librarian, P. O. Box 1663, Los Alamos, New Mexico 87544 (1)
Director, Nuclear Science Center, Louisiana State University, Baton Rouge, Louisiana 70803 (2)
Lovell Foundation for Medical Education and Research, Document Library, 5200 Gibson Blvd., S. E., Albuquerque, New Mexico 87108 (1)
Dr. Ross A. McFarland, Guggenheim Professor of Aerospace Health and Safety, Harvard School of Public Health, 665 Huntington Avenue, Boston, Massachusetts 02115 (1)
Dr. J. I. Marcum, Rand Corporation, 1700 Main Street, Santa Monica, California 90401 (1)
Massachusetts Institute of Technology, M. I. T. Libraries, Technical Reports, Room 14 E-210, Cambridge, Massachusetts 02139 (1)

OTHER (continued)

- Dr. Charles W. Mays, Physics Group Leader, Radiobiology Division, University of Utah, Salt Lake City, Utah 84112 (1)
- Ohio State University, Nuclear Reactor Laboratory, 1298 Kinnear Road, Columbus, Ohio 43212 (1)
- Dr. Harvey M. Patt, Laboratory of Radiobiology, University of California, San Francisco Medical Center, San Francisco, California 94122 (1)
- Purdue University, Nuclear Engineering Library, Lafayette, Indiana 47907 (1)
- Dr. S. M. Reichard, Director, Division of Radiobiology, Medical College of Georgia, Augusta, Georgia 30902 (1)
- University of Rochester, Atomic Energy Project Library, P. O. Box 287, Station 3, Rochester, New York 14620 (1)
- Dr. H. H. Rossi, 630 West 168th Street, New York, N. Y. 10032 (1)
- Dr. Eugene L. Saenger, Director, Radioisotope Laboratory, Cincinnati General Hospital, Cincinnati, Ohio 45229 (1)
- Sandia Corporation Library, P. O. Box 5800, Albuquerque, New Mexico 87115 (1)
- Scientific Committee on the Effects of Atomic Radiation, ATTN: Library, United Nations Room 3267, United Nations Plaza, New York, N. Y. 10017 (1)
- Scope Publications, Franklin Station, P. O. Box 7407, Washington, D. C. 20004 (1)
- Dr. Arthur R. Tamplin, Biophysicist, Information Integration Group, University of California, Lawrence Radiation Laboratory, L-612, Livermore, California 94550 (1)
- Texas A. and M. University, Radiation Biology Laboratory, Texas Engineering Experiment Station, College Station, Texas 77840 (2)
- Texas Nuclear Corporation, ATTN: Director of Research, Box 9267 Allandale Station, Austin, Texas 78756 (1)
- Western Reserve University, Department of Radiology, Division of Radiation Biology, Cleveland, Ohio 44106 (1)
- Mr. Lionel Zamore, 601 Brightwater Court, Brooklyn, New York 11235 (1)

FOREIGN

- International Atomic Energy Agency, Kärltnerring 11, Vienna I, 1010, Austria (1)
- European Atomic Energy Community, C.E.E.A., Library, 51 rue Belliard, Brussels 4, Belgium (1)
- Dr. L. G. Lajtha, Paterson Laboratories, Christie Hospital and Holt Radium Institute, Manchester, England (1)
- Dr. L. F. Lamerton, Biophysics Department, Institute of Cancer Research, Surrey Branch, Belmont, Sutton, Surrey, England (1)
- National Lending Library for Science and Technology, Boston Spa, Yorkshire, England (1)
- Directorate of Medical and Health Services, FAF (Federal Armed Forces), Bonn, Ermekeilstrasse 27, West Germany (1)
- Abteilung für Strahlenbiologie im Institut für Biophysik der Universität Bonn, 53 Bonn-Venusberg, Annaberger Weg 15, Federal Republic of Germany (2)
- Prof. Dr. H. Langendorff, Direktor des Radiologischen Instituts der Universität, 78 Freiburg im Breisgau, Albertstrasse 23, Germany (1)
- Priv.-Doz. Dr. O. Messerschmidt, Radiologisches Institut der Universität, 78 Freiburg im Breisgau, Albertstrasse 23, Germany (1)
- Dr. Helmut Mitschrich, Akademie des Sanitäts- und Gesundheitswesens der Bundeswehr, Spezialstab ATV, 8 München, Schwere Reiterstrasse 4, Germany (2)
- Prof. Dr. F. Wachsmann, Gesellschaft für Strahlenforschung m.b.H., 8042 Neuherberg bei München, Institut für Strahlenschutz, Ingolstadter Landstrasse 1, München, Germany (1)
- Col. Joachim Emde, Direktor, Spezialstab ATV, ABC- und Selbstschuttschule, 8972 Sonthofen 2/Allgäu, Berghoferstrasse 17, West Germany (1)
- Dr. M. Feldman, Section of Cell Biology, The Weizmann Institute of Science, Rehovoth, Israel (1)
- Dr. G. W. Barendsen, Radiobiological Institute TNO, Rijswijk, Netherlands (1)
- Puerto Rico Nuclear Center, ATTN: Reading Room, College Station, Mayaguez, Puerto Rico 00708 (2)
- Dr. H. Cottier, Pathological Institut der Universität, Bern, Switzerland (1)

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Armed Forces Radiobiology Research Institute Defense Nuclear Agency Bethesda, Maryland 20014		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED	
		2b. GROUP N/A	
3. REPORT TITLE ALTERATION OF INTESTINAL VASOACTIVITY DURING THE DEVELOPMENT OF THE GASTROINTESTINAL RADIATION SYNDROME			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
5. AUTHOR(S) (First name, middle initial, last name) J. Kabal, S. J. Baum and L. J. Parkhurst			
6. REPORT DATE September 1971		7a. TOTAL NO. OF PAGES 23	7b. NO. OF REFS 22
8a. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER(S) AFRRI SR71-11	
b. PROJECT NO. NWER XAXM			
c. Task and Subtask C 903		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d. Work Unit 06			
10. DISTRIBUTION STATEMENT Approved for public release; distribution unlimited			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Director Defense Nuclear Agency Washington, D. C. 20305	
13. ABSTRACT <p>The functional vascular integrity of the small intestine in relationship to its morphological alterations during the postirradiation period was examined. Dogs were exposed to 1500 rads (midline tissue dose) of mixed gamma-neutron radiation. Hemodynamic parameters of an <u>in situ</u> intestinal loop were measured in response to the systemic administration of norepinephrine, isoproterenol and gradual bleeding and reinfusion. The results indicate that 48 hours postirradiation when the capillaries are virtually intact and the systemic hemodynamic parameters are not altered yet, the functional integrity of the intestinal vasculature is already significantly deteriorated. The intestinal resistance vessels of the irradiated animals did not exert "autoregulatory escape" as the controls did. After 72 hours the vasocompensation to the sudden blood pressure changes was also markedly changed. The fluctuation of the intestinal resistance values during the bleeding and reinfusion periods demonstrated neurohumoral imbalance in the postirradiation period. On the basis of these results it is postulated that the terminal cardiovascular collapse in irradiated animals might be due to the development of intestinal ischemic shock.</p>			

14.	KEY WORDS	LINK A		LINK B		LINK C	
		ROLE	WT	ROLE	WT	ROLE	WT